

**JANSSEN PHARMACEUTICAL COMPANIES OF JOHNSON & JOHNSON**  
**SELECTED PHARMACEUTICALS IN LATE STAGE U.S. AND E.U. DEVELOPMENT OR REGISTRATION as of 07/15/14**

| Therapeutic Area  | Product Name   | Indication Sought  | U.S. Development Stage  | E.U. Development Stage                               |
|---|--|--|---|--|
| Cardiovascular and Metabolism   | XARELTO® (rivaroxaban)   | Acute Coronary Syndrome (2)<br>Stent Thrombosis for Acute Coronary Syndrome (2)<br>Chronic Heart Failure<br>Prevention of Symptomatic VTE and VTE-related death in high-risk, medically ill patients | FDA Third Complete Response 2/14<br>FDA Second Complete Response 2/14<br>Phase III<br>Phase III |  |
|   | INVOKANA® (canagliflozin)/VOKANAMET® (canagliflozin and metformin fixed dose)  | Fixed Dose Combination with Metformin (IR) (2)   | FDA Complete Response 12/13<br>- Submitted Response to FDA 2/14<br>Phase III                    | Approved 4/14  |
|   | INVOKANA® (canagliflozin)  | Fixed Dose Combination with Metformin (XR) (2)<br>Diabetic nephropathy   | Phase III   | Phase III  |
| Immunology  | STELARA® (ustekinumab)<br>SIMPONI® ARIA™ (golimumab)<br>sirukumab  | Crohn's Disease<br>Rheumatoid Arthritis (IV)<br>Rheumatoid Arthritis   | Phase III<br>Approved 7/13<br>Phase III   | Phase III<br>Application withdrawn 5/14<br>Phase III |
| Infectious Diseases and Vaccines  | PREZISTA® (darunavir) and cobicistat (GS-9350)   | Fixed Dose Combination HIV tablet for treatment naïve patients and treatment experienced patients  | Filed 3/14  | Filed 10/13  |
|   | OLYSIO® (simeprevir)   | Chronic hepatitis C virus (HCV) infection for treatment naïve patients (2)<br>Chronic hepatitis C virus (HCV) infection for treatment experienced patients, relapsers, and non-responders (2)        | Approved 11/13<br>Approved 11/13  | Approved 5/14<br>Approved 5/14                       |
| Neuroscience  | INVEGA® (paliperidone ER OROS)<br>INVEGA® SUSTENNA®/XEPLION® (paliperidone palmitate IM long acting injectable )<br>INVEGA® SUSTENNA® (paliperidone palmitate) | Pediatric indication - adolescent schizophrenia<br>Schizophrenia - 3 month injectable (2)<br><br>Schizoaffective (2)   | Approved 4/11<br>Phase III<br><br>Filed 5/14  | Approved 5/14<br>Phase III                           |
| Oncology  | daratumumab  | Relapsed/refractory multiple myeloma   | Phase III   | Phase III  |
|   | IMBRUVICA® (ibrutinib)   | Relapsed/refractory patients with Chronic Lymphocytic Leukemia (single agent) (randomized study PCYC-1112) (2)   | Filed 4/14  | Phase III  |
|   |  | Relapsed/refractory patients with Chronic Lymphocytic Leukemia (single agent) (single arm phase II study PCYC-1102) (2)  | Approved 2/14   | Filed 10/13  |
|   |  | Relapsed/refractory patients with Chronic Lymphocytic Leukemia (in combination with Bendamustine and Rituximab) (randomized study CLL-3001) (2)  | Phase III   | Phase III  |
|   |  | Treatment naïve patients with Chronic Lymphocytic Leukemia (single agent) (randomized study PCYC-1115) (2)   | Phase III   | Phase III  |
|   |  | Relapsed/refractory with Mantle Cell Lymphoma (single agent) (randomized study MCL-3001) (2)   | Phase III   | Phase III  |
|   |  | Treatment naïve patients with Mantle Cell Lymphoma in combination with Bendamustine and Rituximab (randomized study MCL-3002) (2)  | Phase III   | Phase III  |
|   |  | Relapsed/refractory patients with Mantle Cell Lymphoma (single agent) (single arm phase II study PCYC-1104) (2)  | Approved 11/13  | Filed 10/13  |
|   |  | Relapsed/refractory patients with Indolant Non-Hodgkins Lymphoma in combination with Bendamustine and Rituximab or R-CHOP; (randomized study FLR-3001) (2)   | Phase III   | Phase III  |
|   |  | Newly Diagnosed Non-Germinal Center B-Cell Subtype of Diffuse Large B-Cell Lymphoma in combination with R-CHOP (randomized study DBL3001) (2)  | Phase III   | Phase III  |
| Previously Treated Adults with Waldenstrom's Macroglobulinemia with Rituximab (PCYC-1127) (2) | Phase III  | Phase III  |   |  |
| JNJ56021927/ARN-509   | SYLVANT® (siltuximab)  | Prostate cancer non-metastatic castration-resistant  | Phase III   | Phase III  |
|   |  | Multicentric Castleman's Disease   | Approved 4/14   | Approved 5/14  |
| VELCADE® (bortezomib)   | YONDELIS® (trabectedin)  | Mantle Cell Lymphoma 1st line (2)  | Phase III   | Filed 6/14   |
|   |  | Soft Tissue Sarcoma (2)  | Phase III   |  |
| ZYTIGA® (abiraterone acetate)   |  | Relapsed Ovarian Cancer<br>Prostate Cancer Newly Diagnosed Hormone Naïve Metastatic  | Phase III   | Phase III  |

\* This information is accurate as of the date hereof to the best of the Company's knowledge. Johnson & Johnson assumes no obligation to update this information.

(2) INVOKANA licensed from Mitsubishi Tanabe Pharma Corporation; PREZISTA/cobicistat developed in collaboration with Gilead Sciences; VELCADE developed in collaboration with Millennium: The Takeda Oncology Company; YONDELIS developed in collaboration with PharmaMar S.A.; XARELTO co-developed with Bayer HealthCare; INVEGA SUSTENNA developed in collaboration with Alkermes, Inc.; OLYSIO developed in collaboration with Medivir AB; Daratumumab licensed from Genmab A/S; IMBRUVICA developed in collaboration with Pharmacyclics, Inc.; and Sirukumab developed in collaboration with GlaxoSmithKline.

# Janssen Pharmaceutical Companies of Johnson & Johnson

## Selected Pharmaceutical Pipeline - Recent Approvals/Potential Filings\*

Selective Highlights as of July 15, 2014

| APPROVED 2013/2014  | IN REGISTRATION   | PLANNED FILINGS 2014-2017*  |  |
|---|---|---|--|
| <b>Cardiovascular &amp; Metabolism</b><br><br><b>INVOKANA®</b><br>(canagliflozin)<br>Type 2 diabetes  | <b>Oncology</b><br><br><b>Ibrutinib (EU)</b><br>Previously treated Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma<br>Previously treated Mantle Cell Lymphoma | <b>Immunology</b><br><br><b>Sirukumab</b><br>Rheumatoid arthritis<br><br><b>Guselkumab</b><br>Psoriasis | <b>Oncology</b><br><br><b>Daratumumab</b><br>Relapsed/refractory multiple myeloma<br><br><b>YONDELIS® (US)</b><br>2nd Line soft tissue sarcoma<br><br><b>JNJ56021927 (ARN-509)</b><br>pre-metastatic prostate cancer |
| <b>Infectious Diseases &amp; Vaccines</b><br><br><b>OLYSIO®</b><br>(simeprevir)<br>Chronic Hepatitis C virus<br><br><b>SIRTURO® (EU)</b><br>(bedaquiline)<br>Multi-drug resistant tuberculosis  |   | <b>Neuroscience</b><br><br><b>Esketamine</b><br>Treatment resistant depression                          |  |
| <b>Oncology</b><br><br><b>IMBRUVICA® (US)</b><br>(ibrutinib)<br>Previously treated Mantle Cell Lymphoma<br>Previously treated Chronic Lymphocytic Leukemia<br><br><b>SYLVANT®</b><br>(siltuximab)<br>Multicentric Castleman's disease |   |   |  |

\* Filings/approvals assumed to be in the US and EU unless otherwise noted. This information is accurate as of the date hereof to the best of the Company's knowledge. Johnson & Johnson assumes no obligation to update this information. INVOKANA developed in collaboration with Mitsubishi Tanabe Pharma Corporation; OLYSIO developed in collaboration with Medivir AB; Sirukumab developed in collaboration with GlaxoSmithKline; IMBRUVICA developed in collaboration with Pharmacytics, Inc.; Daratumumab licensed from Genmab A/S; YONDELIS developed in collaboration with Pharma Mar S.A.